



US009409939B2

(12) **United States Patent**
Wang et al.

(10) **Patent No.:** **US 9,409,939 B2**
(45) **Date of Patent:** **Aug. 9, 2016**

(54) **QUANTIFICATION METHOD FOR
REMAINING LIVER FUNCTION AND NOVEL
LIVER RECEPTOR IMAGING AGENT**

(71) Applicant: **INSTITUTE OF NUCLEAR
ENERGY RESEARCH ATOMIC
ENERGY COUNCIL, EXECUTIVE
YUAN**, Taoyuan County (TW)

(72) Inventors: **Mei-Hui Wang**, Taoyuan County (TW);
Wuu-Jyh Lin, Taoyuan County (TW);
Chuan-Yi Chien, Taoyuan County
(TW); **Hung-Man Yu**, Taoyuan County
(TW); **Reiko Takasaka Lee**, Baltimore,
MD (US); **Yuan-Chuan Lee**, Baltimore,
MD (US)

(73) Assignee: **INSTITUTE OF NUCLEAR
ENERGY RESEARCH ATOMIC
ENERGY COUNCIL, EXECUTIVE
YUAN**, Taoyuan (TW)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 155 days.

(21) Appl. No.: **13/858,213**

(22) Filed: **Apr. 8, 2013**

(65) **Prior Publication Data**

US 2013/0225798 A1 Aug. 29, 2013

Related U.S. Application Data

(62) Division of application No. 12/779,374, filed on May
13, 2010, now Pat. No. 8,435,491.

(30) **Foreign Application Priority Data**

Oct. 26, 2009 (TW) 98136146 A

(51) **Int. Cl.**

A61K 51/00 (2006.01)

A61M 36/14 (2006.01)

C07H 15/04 (2006.01)

A61K 51/04 (2006.01)

(52) **U.S. Cl.**

CPC **C07H 15/04** (2013.01); **A61K 51/0491**
(2013.01); **A61K 51/0497** (2013.01)

(58) **Field of Classification Search**

CPC **A61K 51/00**; **A61K 51/08**; **A61K 51/088**;
A61K 51/04; **A61K 51/0491**; **A61K 51/0497**;
C07H 15/04

USPC 424/1.11, 1.65, 1.69, 1.73, 9.1, 9.6;
534/7, 10-16; 514/1, 1.1

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,010,251 A *	3/1977	Green	424/1.49
6,350,431 B1 *	2/2002	Snow	A61K 41/0033 424/9.6
8,142,759 B2 *	3/2012	Wang et al.	424/1.69
8,435,491 B2 *	5/2013	Wang et al.	424/1.69
8,552,163 B2 *	10/2013	Lee et al.	536/17.9
8,961,931 B2 *	2/2015	Duh et al.	424/1.89
9,040,017 B2 *	5/2015	Wang et al.	424/1.11

OTHER PUBLICATIONS

Lee et al, *Bioorganic & Medicinal Chemistry*, 2011, vol. 19, pp.
2494-2500.*

Ljungberg, et al., "3D Absorbed Dose Calculations Based on SPECT:
Evaluation for 111-In/90-Y Therapy using Monte Carlo Simulations.
", 2003, pp. 99-107, vol. 18, No. 1, *Cancer Biotherapy &
Radiopharmaceuticals*.

Royal, et al., "Society of Nuclear Medicine Procedure Guideline for
Hepatic and Splenic Imaging 3.0", Jul. 20, 2003, pp. 53-57, *Society
of Nuclear Medicine Procedure Guidelines Manual*.

Kwon, et al., "Functional Hepatic Volume Measured by Technetium-
99m-Galactosyl-Human Serum Albumin Liver Scintigraphy: Com-
parison Between Hepatocyte Volume and Liver Volume by Com-
puted Tomography", 2001, pp. 541-546, vol. 96, No. 2, *The American
Journal of Gastroenterology*.

Miederer, et al., "Realizing the potential of the Actinium-225
radionuclide generator in targeted alpha particle therapy applica-
tions", 2008, pp. 1371-1382, *Advanced Drug Delivery Reviews*.

Virgolini, et al., "Decreased hepatic function in patients with
hepatoma or liver metastasis monitored by a hepatocyte specific
galactosylated radioligand.", 1990, pp. 937-941, *Macmillan Press
Ltd.*

Korf, et al., "Liver X receptors contribute to the protective immune
response against Mycobacterium tuberculosis in mice", Jun. 2009,
pp. 1826-1837, vol. 119, No. 6, *The Journal of Clinical Investigation*.

Mahley, et al., "Two Independent Lipoprotein Receptors on Hepatic
Membranes of Dog, Swine, and Man", Nov. 1981, pp. 1197-1206,
vol. 68, *The American Society for Clinical Investigation, Inc.*
Yu, et al., "Radiolabeling and MicroSPECT/CT Imaging of a Novel
Multivalent Glycopeptide for Asialoglycoprotein Receptor Imag-
ing", Sep. 2009, vol. 22, *Annals of Nuclear Medicine and Sciences*.

Satoh, et al., "99mTc-GSA liver dynamic SPECT for the preopera-
tive assessment of hepatectomy", 2003, pp. 61-67, vol. 17, No. 1,
Annals of Nuclear Medicine.

* cited by examiner

Primary Examiner — D L Jones

(74) *Attorney, Agent, or Firm* — WPAT, PC; Justin King

(57) **ABSTRACT**

A test indicator for quantifying remaining liver function is
provided. A novel liver receptor imaging agent with liver
targeting property is utilized to develop a method for quanti-
fying remaining liver function to serve as test indicator for
judging the liver failure outcome in clinic, particularly for
judging the necessity of liver transplantation for patients with
liver failure or liver disease. The radioactivity uptake of the
test indicator was negatively correlated with the extent of liver
reserve. The cutoff value of liver reserve for liver transplan-
tation is also disclosed.

1 Claim, 9 Drawing Sheets